

SOME NEW INEQUALITIES OF HERMITE-HADAMARD TYPE FOR h -CONVEX FUNCTIONS ON THE CO-ORDINATES VIA FRACTIONAL INTEGRALS.

ERHAN SET[▲], M. ZEKI SARIKAYA[■], AND HATICE ÖGÜLMÜŞ[♣]

ABSTRACT. By making use of the identity obtained by Sarıkaya, some new Hermite-Hadamard type inequalities for h -convex functions on the co-ordinates via fractional integrals are established. Our results have some relationships with the results of Sarıkaya([16])

1. INTRODUCTION

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$ with $a < b$, then

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}.$$

holds, the double inequality is well known in the literature as Hermite-Hadamard inequality [6]:

Both inequalities hold in the reversed direction if f is concave. For recent results, generalizations and new inequalities related to the Hermite-Hadamard inequality see ([4], [12],[14], [18])

The classical Hermite- Hadamard inequality provides estimates of the mean value of a continuous convex function $f : [a, b] \rightarrow \mathbb{R}$

Let us now consider a bidimensional interval $\Delta := [a, b] \times [c, d]$ in $\mathbb{R}^2$ with $a < b$ and $c < d$. A mapping $f : \Delta \rightarrow \mathbb{R}$ is said to be convex on if the following inequality:

$$f(tx + (1-t)z, ty + (1-t)w) \leq tf(x, y) + (1-t)f(z, w)$$

holds, for all $(x, y), (z, w) \in \Delta$ and $t \in [0, 1]$. If the inequality reversed then f is said to be concave on Δ .

A function $f : \Delta \rightarrow \mathbb{R}$ is said to be on the co-ordinates on Δ if the partial mappings $f_y : [a, b] \rightarrow \mathbb{R}, f_y(u) = f(u, y)$ and $f_x : [c, d] \rightarrow \mathbb{R}, f_x(v) = f(x, v)$ are convex where defined for all $x \in [a, b]$ and $y \in [c, d]$. (see [5])

A formal definition for co-ordinated convex function may be stated as follows:

Definition 1. A function $f : \Delta \rightarrow \mathbb{R}$ will be called co-ordinated convex on Δ , for all $t, k \in [0, 1]$ and $(x, u), (y, w) \in \Delta$, if the following inequality holds:

$$\begin{aligned} & f(tx + (1-t)y, ku + (1-k)w) \\ & \leq tkf(x, u) + k(1-t)f(y, u) + t(1-k)f(x, w) + (1-t)(1-k)f(y, w) \end{aligned}$$

2000 *Mathematics Subject Classification.* 26A33, 26A51, 26D15.

Key words and phrases. Riemann-Liouville fractional integrals, Hermite-Hadamard type inequality, h -convex functions on the co-ordinates.

Clearly, every convex mapping is convex on the co-ordinates, but the converse is not generally true ([5]). Some interesting and important inequalities for convex functions on the co-ordinates can be found in ([9], [10], [17]).

In [1], Alomari and Darus established the following definition of s -convex function in the second sense on co-ordinates.

Definition 2. Consider the bidimensional interval $\Delta := [a, b] \times [c, d]$ in $[0, \infty)^2$ with $a < b$ and $c < d$. The mapping $f : \Delta \rightarrow \mathbb{R}$ is s -convex on Δ if

$$f(\lambda x + (1 - \lambda)z, \lambda y + (1 - \lambda)w) \leq \lambda^s f(x, y) + (1 - \lambda)^s f(z, w)$$

holds for all $(x, y), (z, w) \in \Delta$ with $\lambda \in [0, 1]$, and for some fixed $s \in (0, 1]$.

A function $f : \Delta \rightarrow \mathbb{R}$ is s -convex on Δ is called co-ordinated s -convex on Δ if the partial mappings $f_y : [a, b] \rightarrow \mathbb{R}$, $f_y(u) = f(u, y)$ and $f_x : [c, d] \rightarrow \mathbb{R}$, $f_x(v) = f(x, v)$ are s -convex for all $x \in [a, b]$ and $y \in [c, d]$ with some fixed $s \in (0, 1]$.

In [11], Latif and Alomari give the notion of h -convexity of a function f on a rectangle from the plane $\mathbb{R}^2$ and h -convexity on the co-ordinates on a rectangle from the plane $\mathbb{R}^2$ as follows

Definition 3. Let us consider a bidimensional interval $\Delta := [a, b] \times [c, d]$ in $\mathbb{R}^2$ with $a < b$ and $c < d$. Let $h : J \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a positive function. A mapping $f : \Delta := [a, b] \times [c, d] \rightarrow \mathbb{R}$ is said to be h -convex on Δ , if f is non-negative and if the following inequality:

$$f(\lambda x + (1 - \lambda)z, \lambda y + (1 - \lambda)w) \leq h(\lambda) f(x, y) + h(1 - \lambda) f(z, w)$$

holds, for all $(x, y), (z, w) \in \Delta$ with $\lambda \in (0, 1)$. Let us denote this class of functions by $SX(h, \Delta)$. The function f is said to be h -concave if the inequality reversed. We denote this class of functions by $SV(h, \Delta)$.

A function $f : \Delta \rightarrow \mathbb{R}$ is said to be h -convex on the co-ordinates on Δ if the partial mappings $f_y : [a, b] \rightarrow \mathbb{R}$, $f_y(u) = f(u, y)$ and $f_x : [c, d] \rightarrow \mathbb{R}$, $f_x(v) = f(x, v)$ are h -convex where defined for all $x \in [a, b]$ and $y \in [c, d]$. A formal definition of h -convex functions may also be stated as follows:

Definition 4. [11] A function $f : \Delta \rightarrow \mathbb{R}$ is said to be h -convex on the co-ordinates on Δ , if the following inequality:

(1.1)

$$\begin{aligned} & f(tx + (1 - t)y, ku + (1 - k)w) \\ & \leq h(t)h(k)f(x, u) + h(k)h(1 - t)f(y, u) \\ & + h(t)h(1 - k)f(x, w) + h(1 - t)h(1 - k)f(y, w) \end{aligned}$$

holds for all $t, k \in [0, 1]$ and $(x, u), (x, w), (y, u), (y, w) \in \Delta$.

Obviously, if $h(\alpha) = \alpha$, then all the non-negative convex (concave) functions on Δ belong to the class $SX(h, \Delta)$ ($SV(h, \Delta)$) and if $h(\alpha) = \alpha^s$, where $s \in (0, 1)$, then the class of s -convex on Δ belong to the class $SX(h, \Delta)$. Similarly we can say that if $h(\alpha) = \alpha$, then the class of non-negative convex (concave) functions on the co-ordinates on Δ is contained in the class of h -convex (concave) functions on the co-ordinates on Δ and if $h(\alpha) = \alpha^s$, where $s \in (0, 1)$, then the class of s -convex functions on the co-ordinates on Δ is contained in the class of h -convex functions on the co-ordinates on Δ .

In the following we will give some necessary definitions which are used further in this paper. More details, one can consult [[7], [8], [13]]

Definition 5. Let $f \in L_1[a, b]$. The Riemann-Liouville integrals $J_{a+}^\alpha f$ and $J_{b-}^\alpha f$ order $\alpha > 0$ and $a \geq 0$ are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x < b$$

respectively. Here $\Gamma(\alpha)$ is the Gamma function and $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$.

Definition 6. Let $f \in L_1([a, b] \times [c, d])$. the Riemann-Liouville integrals $J_{a+,c+}^{\alpha,\beta}$, $J_{a+,d-}^{\alpha,\beta}$, $J_{b-,c+}^{\alpha,\beta}$ and $J_{b-,d-}^{\alpha,\beta}$ of order $\alpha, \beta > 0$ with $a, c \geq 0$ are defined by

$$\begin{aligned} & J_{a+,c+}^{\alpha,\beta} f(x, y) \\ &= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_c^y (x-t)^{\alpha-1} (y-s)^{\beta-1} f(t, s) ds dt, \quad x > a, y > c \\ & J_{a+,d-}^{\alpha,\beta} f(x, y) \\ &= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_y^d (x-t)^{\alpha-1} (s-y)^{\beta-1} f(t, s) ds dt, \quad x > a, y < d \\ & J_{b-,c+}^{\alpha,\beta} f(x, y) \\ &= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_c^y (t-x)^{\alpha-1} (y-s)^{\beta-1} f(t, s) ds dt, \quad x < b, y > c \\ & J_{b-,d-}^{\alpha,\beta} f(x, y) \\ &= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_y^d (t-x)^{\alpha-1} (s-y)^{\beta-1} f(t, s) ds dt, \quad x < b, y < d \end{aligned}$$

respectively. Here, Γ is the Gama function,

$$J_{a+,c+}^{0,0} f(x, y) = J_{a+,d-}^{0,0} f(x, y) = J_{b-,c+}^{0,0} f(x, y) = J_{b-,d-}^{0,0} f(x, y) = f(x, y)$$

and

$$J_{a+,c+}^{1,1} f(x, y) = \int_a^x \int_c^y f(t, s) ds dt.$$

For some recent results connected with fractional integral inequalities see ([2], [3], [15], [19]).

In ([16]), Sarıkaya establish the following inequalities of Hadamard's type for co-ordinated convex mapping on a rectangle from the plane $\mathbb{R}^2$:

Theorem 1. Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be co-ordinated convex on $\Delta := [a, b] \times [c, d]$ in $\mathbb{R}^2$ with $0 \leq a < b$, $0 \leq c < d$ and $f \in L_1(\Delta)$. Then one has the inequalities:

(1.2)

$$\begin{aligned}
 & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
 & \leq \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \\
 & \quad \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] \\
 & \leq \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4}.
 \end{aligned}$$

Theorem 2. Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a partial differentiable mapping on $\Delta := [a, b] \times [c, d]$ in $\mathbb{R}^2$ with $0 \leq a < b$, $0 \leq c < d$. If $\left| \frac{\partial^2 f}{\partial t \partial k} \right|$ is a convex function on the co-ordinates on Δ , then one has the inequalities:

(1.3)

$$\begin{aligned}
 & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & \quad + \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \\
 & \quad \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] - A \Big| \\
 & \leq \frac{(b-a)(d-c)}{4(\alpha+1)(\beta+1)} \\
 & \quad \times \left(\left| \frac{\partial^2 f}{\partial k \partial t}(a, c) \right| + \left| \frac{\partial^2 f}{\partial k \partial t}(a, d) \right| + \left| \frac{\partial^2 f}{\partial k \partial t}(b, c) \right| + \left| \frac{\partial^2 f}{\partial k \partial t}(b, d) \right| \right)
 \end{aligned}$$

where

$$\begin{aligned}
 A &= \frac{\Gamma(\beta+1)}{4(d-c)^\beta} \left[J_{c^+}^\beta f(a, d) + J_{c^+}^\beta f(b, d) + J_{d^-}^\beta f(a, c) + J_{d^-}^\beta f(b, c) \right] \\
 & \quad + \frac{\Gamma(\alpha+1)}{4(b-a)^\alpha} \left[J_{a^+}^\alpha f(b, c) + J_{a^+}^\alpha f(b, d) + J_{b^-}^\alpha f(a, c) + J_{b^-}^\alpha f(a, d) \right].
 \end{aligned}$$

Theorem 3. Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a partial differentiable mapping on $\Delta := [a, b] \times [c, d]$ in $\mathbb{R}^2$ with $0 \leq a < b$, $0 \leq c < d$. If $\left| \frac{\partial^2 f}{\partial t \partial k} \right|^q$, $q > 1$, is a convex function

on the co-ordinates on Δ , then one has the inequalities:

$$\begin{aligned}
 (1.4) \quad & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & + \left\{ \frac{\Gamma(\alpha + 1)\Gamma(\beta + 1)}{4(b-a)^\alpha(d-c)^\beta} \right. \\
 & \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] \Big\} - A \\
 & \leq \frac{(b-a)(d-c)}{[(\alpha p + 1)(\beta p + 1)]^{\frac{1}{p}}} \left(\frac{1}{4} \right)^{\frac{1}{q}} \\
 & \times \left(\left| \frac{\partial^2 f}{\partial k \partial t}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial k \partial t}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial k \partial t}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial k \partial t}(b, d) \right|^q \right)^{\frac{1}{q}}
 \end{aligned}$$

where

$$\begin{aligned}
 A = & \frac{\Gamma(\beta + 1)}{4(d-c)^\beta} \left[J_{c^+}^\beta f(a, d) + J_{c^+}^\beta f(b, d) + J_{d^-}^\beta f(a, c) + J_{d^-}^\beta f(b, c) \right] \\
 & + \frac{\Gamma(\alpha + 1)}{4(b-a)^\alpha} \left[J_{a^+}^\alpha f(b, c) + J_{a^+}^\alpha f(b, d) + J_{b^-}^\alpha f(a, c) + J_{b^-}^\alpha f(a, d) \right]
 \end{aligned}$$

and $\frac{1}{p} + \frac{1}{q} = 1$.

In order to prove our main results we need the following lemma (see [16]).

Lemma 1. *Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a partial differentiable mapping on $\Delta := [a, b] \times [c, d]$ in $\mathbb{R}^2$ with $0 \leq a < b$, $0 \leq c < d$. If $\frac{\partial^2 f}{\partial t \partial k} \in L(\Delta)$, then the following equality holds:*

$$\begin{aligned}
 (1.5) \quad & \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \\
 & + \left\{ \frac{\Gamma(\alpha + 1)\Gamma(\beta + 1)}{4(b-a)^\alpha(d-c)^\beta} \right. \\
 & \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] \Big\} \\
 & - \frac{\Gamma(\beta + 1)}{4(d-c)^\beta} \left[J_{c^+}^\beta f(a, d) + J_{c^+}^\beta f(b, d) + J_{d^-}^\beta f(a, c) + J_{d^-}^\beta f(b, c) \right] \\
 & - \frac{\Gamma(\alpha + 1)}{4(b-a)^\alpha} \left[J_{a^+}^\alpha f(b, c) + J_{a^+}^\alpha f(b, d) + J_{b^-}^\alpha f(a, c) + J_{b^-}^\alpha f(a, d) \right]
 \end{aligned}$$

$$\begin{aligned}
&= \frac{(b-a)(d-c)}{4} \left\{ \int_0^1 \int_0^1 t^\alpha k^\beta \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) dk dt \right. \\
&\quad - \int_0^1 \int_0^1 (1-t)^\alpha k^\beta \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) dk dt \\
&\quad - \int_0^1 \int_0^1 t^\alpha (1-k)^\beta \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) dk dt \\
&\quad \left. - \int_0^1 \int_0^1 (1-t)^\alpha (1-k)^\beta \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) dk dt \right\}.
\end{aligned}$$

2. MAIN RESULTS

Theorem 4. Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be h -convex function on the co-ordinates on $\Delta := [a, b] \times [c, d]$ in $\mathbb{R}^2$ and $f \in L_2(\Delta)$. The one has the inequalities:

(2.1)

$$\begin{aligned}
&f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
&\leq \left[h\left(\frac{1}{2}\right)\right]^2 \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{(b-a)^\alpha(d-c)^\beta} \\
&\quad \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c)\right] \\
&\leq \left[h\left(\frac{1}{2}\right)\right]^2 \alpha\beta [f(a, c) + f(a, d) + f(b, c) + f(b, d)] \\
&\quad \times \left[\int_0^1 \int_0^1 t^{\alpha-1} k^{\alpha-1} \right. \\
&\quad \left. \times [h(t)h(k) + h(t)h(1-k) + h(1-t)h(k) + h(1-t)h(1-k)] dk dt\right].
\end{aligned}$$

Proof. According to (1.1) with $x = t_1 a + (1-t_1)b, y = (1-t_1)a + t_1 b, u = k_1 c + (1-k_1)d, w = (1-k_1)c + k_1 d$ and $t = k = \frac{1}{2}$, we find that

(2.2)

$$\begin{aligned}
&f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
&\leq \left[h\left(\frac{1}{2}\right)\right]^2 \\
&\quad \times [f(t_1 a + (1-t_1)b, k_1 c + (1-k_1)d) + f(t_1 a + (1-t_1)b, (1-k_1)c + k_1 d) \\
&\quad + f((1-t_1)a + t_1 b, k_1 c + (1-k_1)d) + f((1-t_1)a + t_1 b, (1-k_1)c + k_1 d)].
\end{aligned}$$

Thus, multiplying both sides of (2.2) by $t_1^{\alpha-1}k_1^{\beta-1}$, then by integrating with respect to (t_1, k_1) on $[0, 1] \times [0, 1]$, we obtain

$$\begin{aligned} & \frac{1}{\alpha\beta} f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\ & \leq \left[h\left(\frac{1}{2}\right) \right]^2 \\ & \quad \times \left[\int_0^1 \int_0^1 t_1^{\alpha-1} k_1^{\beta-1} \right. \\ & \quad \times [f(t_1 a + (1-t_1)b, k_1 c + (1-k_1)d) \\ & \quad + f(t_1 a + (1-t_1)b, (1-k_1)c + k_1 d) \\ & \quad + f((1-t_1)a + t_1 b, k_1 c + (1-k_1)d) \\ & \quad \left. + f((1-t_1)a + t_1 b, (1-k_1)c + k_1 d)] dk_1 ds_1 \right]. \end{aligned}$$

Using the change of the variable, we get

$$\begin{aligned} & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\ & \leq \left[h\left(\frac{1}{2}\right) \right]^2 \frac{\alpha\beta}{(b-a)^\alpha (d-c)^\beta} \left\{ \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} f(x, y) dy dx \right. \\ & \quad + \int_a^b \int_c^d (b-x)^{\alpha-1} (y-c)^{\beta-1} f(x, y) dy dx \\ & \quad + \int_a^b \int_c^d (x-a)^{\alpha-1} (d-y)^{\beta-1} f(x, y) dy dx \\ & \quad \left. + \int_a^b \int_c^d (x-a)^{\alpha-1} (y-c)^{\beta-1} f(x, y) dy dx \right\} \end{aligned}$$

which the first inequality is proved.

For the proof of second inequality (2.1), we first note that if f is a h -convex function on Δ , then, by using (1.1) with $x = a, y = b, u = c, w = d$, it yields

$$\begin{aligned} & f(ta + (1-t)b, kc + (1-k)d) \\ & \leq h(t)h(k)f(a, c) + h(t)h(1-k)f(a, d) \\ & \quad + h(1-t)h(k)f(b, c) + h(1-t)h(1-k)f(b, d) \\ & f((1-t)a + tb, kc + (1-k)d) \\ & \leq h(1-t)h(k)f(a, c) + h(1-t)h(1-k)f(a, d) \\ & \quad + h(t)h(k)f(b, c) + h(t)h(1-k)f(b, d) \\ & f(ta + (1-t)b, (1-k)c + kd) \\ & \leq h(t)h(1-k)f(a, c) + h(t)h(k)f(a, d) \\ & \quad + h(1-t)h(1-k)f(b, c) + h(1-t)h(k)f(b, d) \end{aligned}$$

$$\begin{aligned}
& f((1-t)a + tb, (1-k)c + kd) \\
\leq & h(1-t)h(1-k)f(a, c) + h(1-t)h(k)f(a, d) \\
& + h(t)h(1-k)f(b, c) + h(t)h(k)f(b, d).
\end{aligned}$$

By adding these inequalities we have

(2.3)

$$\begin{aligned}
& f(ta + (1-t)b, kc + (1-k)d) + f((1-t)a + tb, kc + (1-k)d) \\
& + f(ta + (1-t)b, (1-k)c + kd) + f((1-t)a + tb, (1-k)c + kd) \\
\leq & [f(a, c) + f(b, c) + f(a, d) + f(b, d)] \\
& \times [h(t)h(k) + h(t)h(1-k) + h(1-t)h(k) + h(1-t)h(1-k)].
\end{aligned}$$

Then, multiplying both sides of (2.3) by $t^{\alpha-1}k^{\beta-1}$ and integrating with respect to (t, k) over $[0, 1] \times [0, 1]$, we get

$$\begin{aligned}
& \int_0^1 \int_0^1 t^{\alpha-1} k^{\beta-1} \\
& \times [f(ta + (1-t)b, kc + (1-k)d) + f((1-t)a + tb, kc + (1-k)d) \\
& + f(ta + (1-t)b, (1-k)c + kd) + f((1-t)a + tb, (1-k)c + kd)] dk dt \\
\leq & [f(a, c) + f(b, c) + f(a, d) + f(b, d)] \\
& \times \left\{ \int_0^1 \int_0^1 t^{\alpha-1} k^{\beta-1} \right. \\
& \times [h(t)h(k) + h(t)h(1-k) + h(1-t)h(k) + h(1-t)h(1-k)] dk dt \Big\}.
\end{aligned}$$

Here, using the change of the variable we have

$$\begin{aligned}
& \left[h\left(\frac{1}{2}\right) \right]^2 \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{(b-a)^\alpha (d-c)^\beta} \\
& \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] \\
\leq & \left[h\left(\frac{1}{2}\right) \right]^2 \alpha\beta [f(a, c) + f(a, d) + f(b, c) + f(b, d)] \\
& \times \left[\int_0^1 \int_0^1 t^{\alpha-1} k^{\alpha-1} \right. \\
& \times [h(t)h(k) + h(t)h(1-k) + h(1-t)h(k) + h(1-t)h(1-k)] dk dt \Big].
\end{aligned}$$

The proof is completed. $\square$

Remark 1. If we take $h(\alpha) = \alpha$ in Theorem 4, then the inequality (2.1) becomes the inequality (1.2) of Theorem 1.

Corollary 1. If we take $h(\alpha) = \alpha^s$ in Theorem 1, we have the following inequality:

$$\begin{aligned}
& f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
& \leq \left(\frac{1}{2}\right)^{2s} \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{(b-a)^\alpha(d-c)^\beta} \\
& \quad \times \left[J_{a^+,c^+}^{\alpha,\beta} f(b,d) + J_{a^+,d^-}^{\alpha,\beta} f(b,c) + J_{b^-,c^+}^{\alpha,\beta} f(a,d) + J_{b^-,d^-}^{\alpha,\beta} f(a,c) \right] \\
& \leq \left(\frac{1}{2}\right)^{2s} \alpha\beta [f(a,c) + f(a,d) + f(b,c) + f(b,d)] \\
& \quad \times \left(\frac{1}{\alpha+s} + B(\alpha, s+1) \right) \left(\frac{1}{\beta+s} + B(\beta, s+1) \right)
\end{aligned}$$

where B is the Beta function,

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt.$$

Theorem 5. Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a partial differentiable mapping on $\Delta := [a, b] \times [c, d]$ in $\mathbb{R}^2$ with $0 \leq a < b$, $0 \leq c < d$. If $\left| \frac{\partial^2 f}{\partial t \partial k} \right|$ is a h -convex function on the co-ordinates on Δ , then one has the inequalities:

$$\begin{aligned}
(2.4) \quad & \left| \frac{f(a,c) + f(a,d) + f(b,c) + f(b,d)}{4} \right. \\
& + \left\{ \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \right. \\
& \times \left[J_{a^+,c^+}^{\alpha,\beta} f(b,d) + J_{a^+,d^-}^{\alpha,\beta} f(b,c) + J_{b^-,c^+}^{\alpha,\beta} f(a,d) + J_{b^-,d^-}^{\alpha,\beta} f(a,c) \right] \Big\} - A \Big| \\
& \leq \frac{(b-a)(d-c)}{4} \\
& \times \left\{ \left| \frac{\partial^2 f}{\partial t \partial k}(a,c) \right| \int_0^1 \int_0^1 (t^\alpha + (1-t)^\alpha) (k^\beta + (1-k)^\beta) h(t) h(k) dk dt \right. \\
& + \left| \frac{\partial^2 f}{\partial t \partial k}(b,c) \right| \int_0^1 \int_0^1 (t^\alpha + (1-t)^\alpha) (k^\beta + (1-k)^\beta) h(1-t) h(k) dk dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial k}(a,d) \right| \int_0^1 \int_0^1 (t^\alpha + (1-t)^\alpha) (k^\beta + (1-k)^\beta) h(t) h(1-k) dk dt \\
& \left. + \left| \frac{\partial^2 f}{\partial t \partial k}(b,d) \right| \int_0^1 \int_0^1 (t^\alpha + (1-t)^\alpha) (k^\beta + (1-k)^\beta) h(1-t) h(1-k) dk dt \right\}
\end{aligned}$$

where

$$\begin{aligned} A &= \frac{\Gamma(\beta+1)}{4(d-c)^\beta} \left[J_{c^+}^\beta f(a, d) + J_{c^+}^\beta f(b, d) + J_{d^-}^\beta f(a, c) + J_{d^-}^\beta f(b, c) \right] \\ &\quad + \frac{\Gamma(\alpha+1)}{4(b-a)^\alpha} \left[J_{a^+}^\alpha f(b, c) + J_{a^+}^\alpha f(b, d) + J_{b^-}^\alpha f(a, c) + J_{b^-}^\alpha f(a, d) \right]. \end{aligned}$$

Proof. From Lemma 1, we have

$$\begin{aligned} &\left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\ &\quad + \left\{ \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \right. \\ &\quad \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] \Big\} - A \Big| \\ &\leq \frac{(b-a)(d-c)}{4} \left\{ \int_0^1 \int_0^1 t^\alpha k^\beta \left| \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) \right| dk dt \right. \\ &\quad + \int_0^1 \int_0^1 (1-t)^\alpha k^\beta \left| \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) \right| dk dt \\ &\quad + \int_0^1 \int_0^1 t^\alpha (1-k)^\beta \left| \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) \right| dk dt \\ &\quad \left. - \int_0^1 \int_0^1 (1-t)^\alpha (1-k)^\beta \left| \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) \right| dk dt \right\}. \end{aligned}$$

Since $\left| \frac{\partial^2 f}{\partial t \partial k} \right|$ is h-convex function on the co-ordinates on Δ , then one has:

$$\begin{aligned} &\left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\ &\quad + \left\{ \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \right. \\ &\quad \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] \Big\} - A \Big| \end{aligned}$$

$$\begin{aligned}
&\leq \frac{(b-a)(d-c)}{4} \\
&\quad \times \left\{ \left| \frac{\partial^2 f}{\partial t \partial k}(a, c) \right| \int_0^1 \int_0^1 (t^\alpha + (1-t)^\alpha) (k^\beta + (1-k)^\beta) h(t) h(k) dk dt \right. \\
&\quad + \left| \frac{\partial^2 f}{\partial t \partial k}(b, c) \right| \int_0^1 \int_0^1 (t^\alpha + (1-t)^\alpha) (k^\beta + (1-k)^\beta) h(1-t) h(k) dk dt \\
&\quad + \left| \frac{\partial^2 f}{\partial t \partial k}(a, d) \right| \int_0^1 \int_0^1 (t^\alpha + (1-t)^\alpha) (k^\beta + (1-k)^\beta) h(t) h(1-k) dk dt \\
&\quad \left. + \left| \frac{\partial^2 f}{\partial t \partial k}(b, d) \right| \int_0^1 \int_0^1 (t^\alpha + (1-t)^\alpha) (k^\beta + (1-k)^\beta) h(1-t) h(1-k) dk dt \right\}.
\end{aligned}$$

The proof is completed. $\square$

Remark 2. If we take $h(\alpha) = \alpha$ in Theorem 5, then the inequality (2.4) becomes the inequality (1.3) of Theorem 2.

Corollary 2. If we take $h(\alpha) = \alpha^s$ in Theorem 5, we have

$$\begin{aligned}
&\left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
&\quad + \left\{ \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \right. \\
&\quad \left. \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] \right\} - A \Big| \\
&\leq \frac{(b-a)(d-c)}{4} \\
&\quad \times \left[\left| \frac{\partial^2 f}{\partial t \partial k}(a, c) \right| + \left| \frac{\partial^2 f}{\partial t \partial k}(b, c) \right| + \left| \frac{\partial^2 f}{\partial t \partial k}(a, d) \right| + \left| \frac{\partial^2 f}{\partial t \partial k}(b, d) \right| \right] \\
&\quad \times \left(\frac{1}{\alpha+s+1} + B(s+1, \alpha+1) \right) \left(\frac{1}{\beta+s+1} + B(s+1, \beta+1) \right)
\end{aligned}$$

where

$$\begin{aligned}
A &= \frac{\Gamma(\beta+1)}{4(d-c)^\beta} \left[J_{c^+}^\beta f(a, d) + J_{c^+}^\beta f(b, d) + J_{d^-}^\beta f(a, c) + J_{d^-}^\beta f(b, c) \right] \\
&\quad + \frac{\Gamma(\alpha+1)}{4(b-a)^\alpha} \left[J_{a^+}^\alpha f(b, c) + J_{a^+}^\alpha f(b, d) + J_{b^-}^\alpha f(a, c) + J_{b^-}^\alpha f(a, d) \right]
\end{aligned}$$

and B is the Beta function,

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt.$$

Theorem 6. Let $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a partial differentiable mapping on $\Delta := [a, b] \times [c, d]$ in $\mathbb{R}^2$ with $0 \leq a < b$, $0 \leq c < d$. If $\left| \frac{\partial^2 f}{\partial t \partial k} \right|^q, q > 1$, is a h -convex

function on the co-ordinates on Δ , then one has the inequalities:

$$\begin{aligned}
 (2.5) \quad & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & + \left\{ \frac{\Gamma(\alpha + 1) \Gamma(\beta + 1)}{4(b - a)^\alpha (d - c)^\beta} \right. \\
 & \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] \Big\} - A \Big| \\
 & \leq \frac{(b - a)(d - c)}{[(\alpha p + 1)(\beta p + 1)]^{\frac{1}{p}}} \\
 & \times \left(\left| \frac{\partial^2 f}{\partial k \partial t}(a, c) \right|^q \int_0^1 \int_0^1 h(t) h(k) dk dt \right. \\
 & + \left| \frac{\partial^2 f}{\partial k \partial t}(a, d) \right|^q \int_0^1 \int_0^1 h(t) h(1 - k) dk dt \\
 & + \left| \frac{\partial^2 f}{\partial k \partial t}(b, c) \right|^q \int_0^1 \int_0^1 h(1 - t) h(k) dk dt \\
 & \left. + \left| \frac{\partial^2 f}{\partial k \partial t}(b, d) \right|^q \int_0^1 \int_0^1 h(1 - t) h(1 - k) dk dt \right)^{\frac{1}{q}}
 \end{aligned}$$

where

$$\begin{aligned}
 A = & \frac{\Gamma(\beta + 1)}{4(d - c)^\beta} \left[J_{c^+}^\beta f(a, d) + J_{c^+}^\beta f(b, d) + J_{d^-}^\beta f(a, c) + J_{d^-}^\beta f(b, c) \right] \\
 & + \frac{\Gamma(\alpha + 1)}{4(b - a)^\alpha} \left[J_{a^+}^\alpha f(b, c) + J_{a^+}^\alpha f(b, d) + J_{b^-}^\alpha f(a, c) + J_{b^-}^\alpha f(a, d) \right]
 \end{aligned}$$

and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. From Lemma 1, we have

$$\begin{aligned}
 & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & + \left\{ \frac{\Gamma(\alpha + 1) \Gamma(\beta + 1)}{4(b - a)^\alpha (d - c)^\beta} \right. \\
 & \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] \Big\} - A \Big|
 \end{aligned}$$

$$\begin{aligned}
\leq & \frac{(b-a)(d-c)}{4} \left\{ \int_0^1 \int_0^1 t^\alpha k^\beta \left| \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) \right| dk dt \right. \\
& + \int_0^1 \int_0^1 (1-t)^\alpha k^\beta \left| \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) \right| dk dt \\
& + \int_0^1 \int_0^1 t^\alpha (1-k)^\beta \left| \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) \right| dk dt \\
& \left. - \int_0^1 \int_0^1 (1-t)^\alpha (1-k)^\beta \left| \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) \right| dk dt \right\}.
\end{aligned}$$

By using the well known Hölder's inequality for double integrals, we get

$$\begin{aligned}
& \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
& + \left\{ \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \right. \\
& \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^+}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^+}^{\alpha, \beta} f(a, c) \right] \Big\} - A \Big| \\
\leq & \frac{(b-a)(d-c)}{4} \left\{ \left(\int_0^1 \int_0^1 t^{p\alpha} k^{p\beta} dk dt \right)^{\frac{1}{p}} + \left(\int_0^1 \int_0^1 (1-t)^{p\alpha} k^{p\beta} dk dt \right)^{\frac{1}{p}} \right. \\
& \left. \left(\int_0^1 \int_0^1 t^{p\alpha} (1-k)^{p\beta} dk dt \right)^{\frac{1}{p}} + \left(\int_0^1 \int_0^1 (1-t)^{p\alpha} (1-k)^{p\beta} dk dt \right)^{\frac{1}{p}} \right\} \\
& \left(\int_0^1 \int_0^1 \left| \frac{\partial^2 f}{\partial t \partial k} (ta + (1-t)b, kc + (1-k)d) \right|^q dk dt \right)^{\frac{1}{q}}
\end{aligned}$$

Since $\left| \frac{\partial^2 f}{\partial t \partial k} \right|^q$ is h-convex function on the co-ordinates on Δ , then one has:

$$\begin{aligned}
& \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
& + \left\{ \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \right. \\
& \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^+}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^+}^{\alpha, \beta} f(a, c) \right] \Big\} - A \Big|
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{(b-a)(d-c)}{[(\alpha p+1)(\beta p+1)]^{\frac{1}{p}}} \\
&\quad \times \left(\left| \frac{\partial^2 f}{\partial k \partial t}(a, c) \right|^q \int_0^1 \int_0^1 h(t) h(k) dk dt \right. \\
&\quad + \left| \frac{\partial^2 f}{\partial k \partial t}(a, d) \right|^q \int_0^1 \int_0^1 h(t) h(1-k) dk dt \\
&\quad + \left| \frac{\partial^2 f}{\partial k \partial t}(b, c) \right|^q \int_0^1 \int_0^1 h(1-t) h(k) dk dt \\
&\quad \left. + \left| \frac{\partial^2 f}{\partial k \partial t}(b, d) \right|^q \int_0^1 \int_0^1 h(1-t) h(1-k) dk dt \right)^{\frac{1}{q}}
\end{aligned}$$

and the proof is completed. $\square$

Remark 3. If we take $h(\alpha) = \alpha$ in Theorem 6, then the inequality (2.5) becomes the inequality (1.4) of Theorem 3.

Corollary 3. If we take $h(\alpha) = \alpha^s$ in Theorem 5, we have

$$\begin{aligned}
&\left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
&\quad + \left\{ \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \right. \\
&\quad \times \left[J_{a^+, c^+}^{\alpha, \beta} f(b, d) + J_{a^+, d^-}^{\alpha, \beta} f(b, c) + J_{b^-, c^+}^{\alpha, \beta} f(a, d) + J_{b^-, d^-}^{\alpha, \beta} f(a, c) \right] \Big\} - A \Big| \\
&\leq \frac{(b-a)(d-c)}{[(\alpha p+1)(\beta p+1)]^{\frac{1}{p}}} \\
&\quad \times \left(\frac{1}{(s+1)^2} \left[\left| \frac{\partial^2 f}{\partial k \partial t}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial k \partial t}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial k \partial t}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial k \partial t}(b, d) \right|^q \right] \right)^{\frac{1}{q}}
\end{aligned}$$

where

$$\begin{aligned}
A &= \frac{\Gamma(\beta+1)}{4(d-c)^\beta} \left[J_{c^+}^\beta f(a, d) + J_{c^+}^\beta f(b, d) + J_{d^-}^\beta f(a, c) + J_{d^-}^\beta f(b, c) \right] \\
&\quad + \frac{\Gamma(\alpha+1)}{4(b-a)^\alpha} \left[J_{a^+}^\alpha f(b, c) + J_{a^+}^\alpha f(b, d) + J_{b^-}^\alpha f(a, c) + J_{b^-}^\alpha f(a, d) \right].
\end{aligned}$$

REFERENCES

- [1] M. Alomari and M. Darus, The Hadamard's inequality for s -convex function of 2-variables on the co-ordinates, Int. Journal of Math. Analysis, 2 (13) (2008), 629-638.
- [2] S. Belarbi and Z. Dahmani, On some new fractional integral inequalities, J. Ineq. Pure and Appl. Math., 10(3) (2009), Art. 86.
- [3] Z. Dahmani, L. Tabharit, S. Taf, Some fractional integral inequalities, Nonl. Sci. Lett. A, 1(2) (2010), 155-160.
- [4] S.S. Dragomir, On some new inequalities of Hermite-Hadamard type for m -convex functions, Tamkang J. Math., 3 (1) (2002).

- [5] S.S. Dragomir, On Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane, Taiwanese J. Math. 4 (2001), 775–788.
- [6] S.S. Dragomir, C.E.M. Pearce, Selected topics on Hermite–Hadamard inequalities and applications, RGMIA Monographs, Victoria University, 2000.
- [7] R. Gorenflo, F. Mainardi, Fractional calculus: integral and differential equations of fractional order, Springer Verlag, Wien (1997), 223–276.
- [8] S. Miller and B. Ross, An introduction to the fractional calculus and fractional differential equations, John Wiley & Sons, USA, 1993, p.2.
- [9] M.A. Latif, M. Alomari, Hadamard-type inequalities for product two convex functions on the co-ordinates. Int Math Forum. 4(47), 2327–2338 (2009).
- [10] M.A. Latif and S. Hussain, New inequalities of Ostrowski type for co-ordinated convex functions via fractional integrals, Journal of Fractional Calculus and Appl. 2(9) (2012), 1–15.
- [11] M.A. Latif, On Hadamard-type inequalities for h-convex functions on the co-ordinates, Int. Journal of Math. Analysis, 3 (33) (2009), 1645 – 1656.
- [12] C.E.M. Pearce, J. Pečarić, Inequalities for differentiable mappings with application to special means and quadrature formula, Appl. Math. Lett. 13 (2000), 51—55.
- [13] I. Podlubni, Fractional differential equations, Academic Press, San Diego, 1999.
- [14] M. Z Sarikaya., A. Saglam and H. Yildirim, On some Hadamard type inequalities for h-convex functions, Journal Math. Ineq. 2(3) (2008), 335–341.
- [15] M.Z. Sarikaya and H. Yaldiz, On weighted Montgomery identities for Riemann- Liouville fractional integrals, Konuralp Journal of Math., 1(1) (2013), 48–53.
- [16] M.Z. Sarikaya, On the Hermite-Hadamard-type inequalities for co-ordinated convex function via fractional integrals (submitted).
- [17] M.Z. Sarikaya, E. Set, M.E. Özdemir, S.S. Dragomir, New some Hadamard's type inequalities for co-ordinated convex functions. Tamsui Oxford Journal of Information and Mathematical Sciences, 28(2) (2012), 137–152.
- [18] E. Set, M.E. Özdemir, S.S. Dragomir, On the Hermite–Hadamard inequality and other integral inequalities involving two functions, J. Inequal. Appl. (2010), p. 9 Article ID 148102.
- [19] E. Set, New inequalities of Ostrowski type for mappings whose derivatives are s-convex in the second sense via fractional integrals, Comput. Math. Appl., 63 (2012), 1147–1154.

▲DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, ORDU UNIVERSITY, ORDU-TURKEY

E-mail address: erhanset@yahoo.com

■DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY

E-mail address: sarikayamz@gmail.com

♣DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY

E-mail address: haticeogulmus@hotmail.com